

# The structure of graphs with a vital linkage of order 2\*

Dillon Mayhew<sup>†</sup>   Geoff Whittle<sup>†</sup>   Stefan H. M. van Zwam<sup>‡</sup>

July 11, 2011

## Abstract

A *linkage of order  $k$*  of a graph  $G$  is a subgraph with  $k$  components, each of which is a path. A linkage is *vital* if it spans all vertices, and no other linkage connects the same pairs of end vertices. We give a characterization of the graphs with a vital linkage of order 2: they are certain minors of a family of highly structured graphs.

## 1 Introduction

Robertson and Seymour [4] defined a *linkage* in a graph  $G$  as a subgraph in which each component is a path. The *order* of a linkage is the number of components. A linkage  $L$  of order  $k$  is *unique* if no other collection of paths connects the same pairs of vertices, it is *spanning* if  $V(L) = V(G)$ , and it is *vital* if it is both unique and spanning. Graphs with a vital linkage are well-behaved. For instance, Robertson and Seymour proved the following:

**Theorem 1.1** (Robertson and Seymour [4, Theorem 1.1]). *There exists an integer  $w$ , depending only on  $k$ , such that every graph with a vital linkage of order  $k$  has tree width at most  $w$ .*

Note that Robertson and Seymour use the term  $p$ -linkage to denote a linkage with  $p$  terminals. Robertson and Seymour's proof of this theorem is surprisingly elaborate, and uses their structural description of graphs with no large clique-minor. Recently Kawarabayashi and Wollan

---

\*The research of all authors was partially supported by a grant from the Marsden Fund of New Zealand. The first author was also supported by a FRST Science & Technology post-doctoral fellowship. The third author was also supported by the Netherlands Organization for Scientific Research (NWO).

<sup>†</sup>School of Mathematics, Statistics and Operations Research, Victoria University of Wellington, New Zealand. E-mail: [Dillon.Mayhew@msor.vuw.ac.nz](mailto:Dillon.Mayhew@msor.vuw.ac.nz), [Geoff.Whittle@msor.vuw.ac.nz](mailto:Geoff.Whittle@msor.vuw.ac.nz)

<sup>‡</sup>Centrum Wiskunde en Informatica, Postbus 94079, 1090 GB Amsterdam, The Netherlands. E-mail: [Stefan.van.Zwam@cw.nl](mailto:Stefan.van.Zwam@cw.nl)




Figure 1: The graph  $K_{2,4}$ .




Figure 2: The graphs  $\ddot{U}_4$  and  $\ddot{U}_5$ .

[2] proved a strengthening of this result. Their shorter proof avoids using the structure theorem.

Our interest in linkages, in particular those of order 2, stems from quite a different area of research: matroid theory. Truemper [5] studied a class of binary matroids that he calls *almost regular*. His proofs lean heavily on a class of matroids that are single-element extensions of the cycle matroids of graphs with a vital linkage of order 2. These matroids turned up again in the excluded-minor characterization of matroids that are either binary or ternary, by Mayhew et al. [3].

Truemper proves that an almost regular matroid can be built from one of two specific matroids by certain  $\Delta - Y$  operations. This is a deep result, but it does not yield bounds on the branch width of these matroids. In a forthcoming paper the authors of this paper, together with Chun, will give an explicit structural description of the class of almost regular matroids [1]. The main result of this paper will be of use in that project.

To state our main result we need a few more definitions. Fix a graph  $G$  and a spanning linkage  $L$  of order  $k$ . A *path edge* is a member of  $E(L)$ ; an edge in  $E(G) - E(L)$  is called a *chord* if its endpoints lie in a single path, and a *rung edge* otherwise. If  $L$  is vital, then  $G$  cannot have any chords.

A *linkage minor* of  $G$  with respect to a (chordless) linkage  $L$  is a minor  $H$  of  $G$  such that all path edges in  $E(G) - E(H)$  have been contracted, and all rung edges in  $E(G) - E(H)$  have been deleted. If the linkage  $L$  is clear from the context we simply say that  $H$  is a linkage minor of  $G$ . Moreover, let  $G$  be a graph with a chordless 2-linkage  $L$ . If  $G$  has a linkage minor isomorphic to  $K_{2,4}$ , such that the terminals of  $L$  are mapped to the degree-2 vertices of  $K_{2,4}$ , we say that  $G$  has an *XX linkage minor* (cf. Figure 1).


For each integer  $n$ , the graph  $\ddot{U}_n$  is the graph with  $V(\ddot{U}_n) = \{v_1, \dots, v_n\} \cup \{u_1, \dots, u_n\}$ , and

$$E(\ddot{U}_n) = \{v_i v_{i+1} \mid i = 1, \dots, n-1\} \cup \{u_i u_{i+1} \mid i = 1, \dots, n-1\} \cup \{u_i v_i \mid i = 1, \dots, n\} \cup \{u_i v_{n+1-i} \mid i = 1, \dots, n\}. \quad (1)$$

We denote by  $L_n$  the linkage of  $\ddot{U}_n$  consisting of all edges  $v_i v_{i+1}$  and  $u_i u_{i+1}$  for  $i = 1, \dots, n-1$ . In Figure 2 the graphs  $\ddot{U}_4$  and  $\ddot{U}_5$  are depicted.

Finally, we say that  $G$  is a *Truemper graph* if  $G$  is a linkage minor of  $\ddot{U}_n$  for some  $n$ . The main result of this paper is the following:

**Theorem 1.2.** *Let  $G$  be a graph. The following statements are equivalent:*



**Figure 3:** The graph  $\ddot{U}_6$ . The linkage is formed by the two diagonally drawn paths.

- (i)  $G$  has a vital linkage of order 2;
- (ii)  $G$  has a chordless spanning linkage of order 2 with no  $XX$  linkage minor;
- (iii)  $G$  is a Truemper graph.

Robertson and Seymour [4] commented, without proof, that graphs with a vital linkage with  $k \leq 5$  terminal vertices have path width at most  $k$ . A weaker claim is the following:

**Corollary 1.3.** *Let  $G$  be a graph with a vital linkage of order 2. Then  $G$  has path width at most 4.*

Another consequence of our result is that graphs with a vital linkage of order 2 embed in the projective plane:

**Corollary 1.4.** *Let  $G$  be a graph with a vital linkage of order 2. Then  $G$  can be embedded on a Möbius strip.*


Both corollaries can be seen to be true by considering an alternative depiction of  $\ddot{U}_{2n}$ , analogous to Figure 3.

## 2 Proof of Theorem 1.2

We start with a few more definitions. Suppose  $L$  is a linkage of order 2 with components  $P_1$  and  $P_2$ , such that the terminal vertices of  $P_1$  are  $s_1$  and  $t_1$ , and those of  $P_2$  are  $s_2$  and  $t_2$ . We order the vertices on the paths in a natural way, as follows. If  $v$  and  $w$  are vertices of  $P_i$ , then we say that  $v$  is (strictly) to the left of  $w$  if the graph distance from  $s_i$  to  $v$  in the subgraph  $P_i$  is (strictly) smaller than the graph distance from  $s_i$  to  $w$ . The notion to the right is defined analogously.

We will frequently use the following elementary observation, whose proof we omit.

**Lemma 2.1.** *Let  $G$  be a graph with a chordless spanning linkage  $L$  of order 2. Let  $P_1$  and  $P_2$  be the components of  $L$ , with terminal vertices respectively  $s_1, t_1$  and  $s_2, t_2$ . Let  $H$  be a linkage minor of  $G$ . If  $v$  and  $w$  are on  $P_i$ , and  $v$  is to the left of  $w$ , then the vertex corresponding to  $v$  in  $H$  is to the left of the vertex corresponding to  $w$  in  $H$ .*



**Figure 4:** Detail of the proof of Lemma 2.2.

Without further ado we dive into the proof, which will consist of a sequence of lemmas. The first deals with the equivalence of the first two statements in the theorem.

**Lemma 2.2.** *Let  $G$  be a graph with a chordless spanning linkage  $L$  of order 2. Then  $L$  is vital if and only if  $G$  has no  $XX$  linkage minor with respect to  $L$ .*

*Proof.* First we suppose that there exists a graph  $G$  with a non-vital chordless spanning linkage  $L$  of order 2 such that  $G$  has no  $XX$  linkage minor. Let  $P_1, P_2$  be the paths of  $L$ , where  $P_1$  runs from  $s_1$  to  $t_1$ , and  $P_2$  runs from  $s_2$  to  $t_2$ . Let  $P'_1, P'_2$  be different paths connecting the same pairs of vertices. Without loss of generality,  $P'_1 \neq P_1$ . But then  $P'_1$  must meet  $P_2$ , so  $P'_2 \neq P_2$ . Let  $e = v_1 v_2$  be an edge of  $P'_1$  such that the subpath  $s_1 - v_1$  of  $P'_1$  is also a subpath of  $P_1$ , but  $e$  is not an edge of  $P_1$ . Let  $f = u_2 u_1$  be an edge of  $P'_1$  such that the subpath  $u_1 - t_1$  of  $P'_1$  is also a subpath of  $P_2$ , but  $f$  is not an edge of  $P_2$ . Similarly, let  $e' = v_2' v_1'$  be an edge of  $P'_2$  such that the subpath  $s_2 - v_2'$  of  $P'_2$  is also a subpath of  $P_2$ , but  $e'$  is not an edge of  $P_2$ . Let  $f' = u_1' u_2'$  be an edge of  $P'_2$  such that the subpath  $u_2' - t_2$  of  $P'_2$  is also a subpath of  $P_2$ , but  $f'$  is not on  $P_2$ . See Figure 4.

Since  $P'_1$  and  $P'_2$  are vertex-disjoint,  $v_2'$  must be strictly to the left of  $v_2$  and  $u_2$ . For the same reason,  $v_1'$  must be strictly between  $v_1$  and  $u_1$ . Likewise,  $u_2'$  must be strictly to the right of  $v_2$  and  $u_2$ , and  $u_1'$  must be strictly between  $v_1$  and  $u_1$ . Now construct a linkage minor  $H$  of  $G$ , as follows. Contract all edges on the subpaths  $s_1 - v_1$ ,  $v_1' - u_1'$ , and  $u_1 - t_1$  of  $P_1$ , contract all edges on the subpaths  $s_2 - v_2'$ ,  $v_2 - u_2$ , and  $u_2' - t_2$  of  $P_2$ , delete all rung edges but  $\{e, f, e', f'\}$ , and contract all but one of the edges of each series class in the resulting graph. Clearly  $H$  is isomorphic to  $XX$ , a contradiction.

Conversely, suppose that  $G$  has an  $XX$  linkage minor, but that  $L$  is unique. Clearly having a vital linkage is preserved under taking linkage minors. But  $XX$  has two linkages, a contradiction.  $\square$

Next we show that the third statement of Theorem 1.2 implies the second.

**Lemma 2.3.** *For all  $n$ ,  $\ddot{U}_n$  has no  $XX$  linkage minor with respect to  $L_n$ .*

*Proof.* The result holds for  $n \leq 2$ , because then  $|V(\ddot{U}_n)| < |V(XX)|$ . Suppose the lemma fails for some  $n \geq 3$ , but is valid for all smaller  $n$ . Every edge of  $XX$  is incident with exactly one of the four end vertices of the paths. Hence all rung edges incident with at least two of the four end vertices are not in any  $XX$  linkage minor. But after deleting those edges

from  $\ddot{U}_n$  the end vertices have degree one, and hence the edges incident with them will not be in any  $XX$  linkage minor. Contracting these four edges produces  $\ddot{U}_{n-2}$ , a contradiction.  $\square$

Reversing a path  $P_i$  means exchanging the labels of vertices  $s_i$  and  $t_i$ , thereby reversing the order on the vertices of the path.

**Lemma 2.4.** *Let  $G$  be a graph, and  $L$  a chordless spanning linkage of order 2 of  $G$  consisting of paths  $P_1$ , running from  $s_1$  to  $t_1$ , and  $P_2$ , running from  $s_2$  to  $t_2$ . If  $G$  has no  $XX$  linkage minor, then  $G$  is a linkage minor of  $\ddot{U}_n$  with respect to  $L_n$  for some integer  $n$ , such that  $L$  is a contraction of  $L_n$ .*

*Proof.* Suppose the statement is false. Let  $G$  be a counterexample with as few edges as possible. If some end vertex of a path, say  $s_1$ , has degree one (with  $e = s_1v$  the only edge), then we can embed  $G/e$  in  $\ddot{U}_n$  for some  $n$ . Let  $G'$  be obtained from  $\ddot{U}_n$  by adding four vertices  $s'_1, t'_1, s'_2, t'_2$ , and edges  $s'_1v_1, s'_1s'_2, s'_1t'_2, s'_2u_1, s'_2t'_1, v_1t'_1, u_1t'_2, t'_1t'_2$ . Then  $G'$  is isomorphic to  $\ddot{U}_{n+2}$ , and  $G'$  certainly has  $G$  as linkage minor.

Hence we may assume that each end vertex of  $P_1$  and  $P_2$  has degree at least two. Suppose no rung edge runs between two of these end vertices. Then it is not hard to see that  $G$  has an  $XX$  minor, a contradiction. Therefore some two end vertices must be connected. By reversing paths as necessary, we may assume there is an edge  $e = s_1s_2$ .

By our assumption,  $G \setminus e$  can be embedded in  $\ddot{U}_n$  for some  $n$ . Again, let  $G'$  be obtained from  $\ddot{U}_n$  by adding four vertices  $s'_1, t'_1, s'_2, t'_2$ , and edges  $s'_1v_1, s'_1s'_2, s'_1t'_2, s'_2u_1, s'_2t'_1, v_1t'_1, u_1t'_2, t'_1t'_2$ . Then  $G'$  is isomorphic to  $\ddot{U}_{n+2}$ , and  $G'$  certainly has  $G$  as linkage minor, a contradiction.  $\square$

As an aside, it is possible to prove a stronger version of the previous lemma. We say a partition  $(A, B)$  of the rung edges is *valid* if the edges in  $A$  are pairwise non-crossing, and the edges in  $B$  are pairwise non-crossing after reversing one of the paths. One can show:

- Each Truemper graph has a valid partition.
- For every valid partition  $(A, B)$  of a Truemper graph  $G$ , some  $\ddot{U}_n$  has  $G$  as linkage minor in such a way that  $(A, B)$  extends to a valid partition of  $\ddot{U}_n$ .

Now we have all ingredients of our main result.

*Proof of Theorem 1.2.* From Lemma 2.2 we learn that  $(i) \Leftrightarrow (ii)$ . From Lemma 2.3 we learn that  $(iii) \Rightarrow (ii)$ , and from Lemma 2.4 we conclude that  $(ii) \Rightarrow (iii)$ .  $\square$

## References

- [1] Carolyn Chun, Dillon Mayhew, Geoff Whittle, and Stefan H. M. van Zwam. The structure of binary Fano-fragile matroids. In preparation.
- [2] Ken-ichi Kawarabayashi and Paul Wollan. A shorter proof of the graph minor algorithm: the unique linkage theorem. In *STOC*, pages 687–694, 2010.

- [3] Dillon Mayhew, Bogdan Oporowski, James Oxley, and Geoff Whittle. The excluded minors for the matroids that are binary or ternary. Submitted, 2009.
- [4] Neil Robertson and P. D. Seymour. Graph minors. XXI. graphs with unique linkages. *J. Combin. Theory Ser. B*, 99(3):583–616, 2009.
- [5] K. Truemper. A decomposition theory of matroids. VI. Almost regular matroids. *J. Combin. Theory Ser. B*, 55(2):235–301, 1992. ISSN 0095-8956.